



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Full genome sequence of Escherichia coli K-15KW01, an uropathogenic E. coli B2:ST127 isolate harboring a chromosomally-encoded blaCTX-M-15 gene

Zurfluh, Katrin ; Tasara, Taurai ; Stephan, Roger

Abstract: We present here the full-genome sequence of Escherichia coli K-15KW01, an extended-spectrum-lactamase-producing uropathogenic strain. Assembly and annotation of the draft genome resulted in a 5,154,641-bp chromosome and revealed a chromosomally contained blaCTX-M-15 gene embedded at the right-hand extremity of an ISEcp1 element in a plasmid-like structure (36,907 bp).

DOI: <https://doi.org/10.1128/genomeA.00927-16>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-129956>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Zurfluh, Katrin; Tasara, Taurai; Stephan, Roger (2016). Full genome sequence of Escherichia coli K-15KW01, an uropathogenic E. coli B2:ST127 isolate harboring a chromosomally-encoded blaCTX-M-15 gene. Genome Announcements, 4(5):e00927-16.

DOI: <https://doi.org/10.1128/genomeA.00927-16>

Full-Genome Sequence of *Escherichia coli* K-15KW01, a Uropathogenic *E. coli* B2 Sequence Type 127 Isolate Harboring a Chromosomally Carried *bla*_{CTX-M-15} Gene

Katrin Zurfluh, Taurai Tasara, Roger Stephan

Institute for Food Safety and Hygiene, Vetsuisse Faculty, University of Zurich, Zürich, Switzerland

We present here the full-genome sequence of *Escherichia coli* K-15KW01, an extended-spectrum- β -lactamase-producing uropathogenic strain. Assembly and annotation of the draft genome resulted in a 5,154,641-bp chromosome and revealed a chromosomally contained *bla*_{CTX-M-15} gene embedded at the right-hand extremity of an *ISEcp1* element in a plasmid-like structure (36,907 bp).

Received 9 July 2016 Accepted 18 July 2016 Published 1 September 2016

Citation Zurfluh K, Tasara T, Stephan R. 2016. Full-genome sequence of *Escherichia coli* K-15KW01, a uropathogenic *E. coli* B2 sequence type 127 isolate harboring a chromosomally carried *bla*_{CTX-M-15} gene. *Genome Announc* 4(5):e00927-16. doi:10.1128/genomeA.00927-16.

Copyright © 2016 Zurfluh et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Roger Stephan, stephanr@fsafety.uzh.ch.

The production of extended-spectrum β -lactamases (ESBLs) is one of the most important mechanisms of antibacterial resistance in *Enterobacteriaceae*. Most ESBLs can be divided into 4 groups: TEM, SHV, OXA, and CTX-M types (1). Currently, CTX-Ms are the most prevalent type of ESBLs described (2, 3). The last decade has seen a high degree of diversification of these enzymes and an explosive global spread driven primarily by their carriage on resistance plasmids and by the spread of extraintestinal pathogenic *Escherichia coli* clones (4). Meanwhile, it has become widely recognized that the dissemination of ESBL-producing bacteria is an issue that is no longer restricted to the medical health care system. In the last few years, there has been an increase in the detection of ESBL-producing strains in the general community (5, 6).

In a recent study, we isolated an ESBL-producing *E. coli* isolate (B2 sequence type 127 [ST127]) harboring a *bla*_{CTX-M-15} gene from a healthy asymptomatic carrier shedding the same strain for 5 months in fecal samples (7). Despite repeated attempts, the *bla*_{CTX-M-15} gene from this isolate could not be transferred by conjugation. Southern blot hybridization provided evidence for a chromosomal location of *bla*_{CTX-M-15} (data not shown). Therefore, genomic DNA was isolated from K-15KW01 and subjected to sequencing using Pacific Biosciences single-molecule real-time (SMRT) technology at the ChunLab at Seoul National University, South Korea. The K-15KW01 genome was assembled *de novo* using the SMRT Analysis 2.3.0 software to a single chromosome of 5,154,641 bp in size, with a G+C content of 50.4%. Genome annotation was conducted through the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (http://www.ncbi.nlm.nih.gov/genome/annotation_prok/) (8).

By full-genome sequencing, the chromosomal location of the *bla*_{CTX-M-15} gene could be proven. The *bla*_{CTX-M-15} gene was embedded at the right-hand extremity of an *ISEcp1* element in a plasmid-like structure (36,907 bp) located between a putative gene encoding for an oxalyl-coenzyme A (CoA) decarboxylase

domain (positions 978270 to 978743) and a formyl-CoA-oxalate CoA-transferase (gene *frc*) (positions 1015704 to 1016954). Moreover, on this plasmid-like structure, a tunicamycin resistance determinant (*tmrB*), an aminoglycoside N(3')-acetyl transferase III (*aacC2*), the *cr* variant of *aac(6')-Ib* encoding an aminoglycoside acetyltransferase conferring resistance to fluoroquinolones, the β -lactamase gene *bla*_{OXA-1}, and the chloramphenicol acetyltransferase *catB3* were carried.

E. coli B2:ST127 has recently been described as an emerging clone with extremely high uropathogenic potential (9) and has been associated with community-acquired urinary tract infections (CAUTI) in Britain (10). The detection of uropathogenic *E. coli* (UPEC) B2:ST127 isolates harboring chromosomally carried *bla*_{CTX-M-15} in a healthy person is a matter of concern, since UPEC isolates spread easily via person-to-person contact or fecal-oral transmission (11).

Accession number(s). Sequence and annotation data of the genome of *E. coli* strain K-15KW01 were deposited in the GenBank database with the accession no. [CP016358](https://doi.org/10.1101/16358).

FUNDING INFORMATION

This work was supported by funding of the University of Zurich and the Swiss Federal Office of Public Health to Katrin Zurfluh.

REFERENCES

1. Bush K, Jacoby GA. 2010. Updated functional classification of beta-lactamases. *Antimicrob Agents Chemother* 54:969–976. <http://dx.doi.org/10.1128/AAC.01009-09>.
2. Cantón R, González-Alba JM, Galán JC. 2012. CTX-M enzymes: origin and diffusion. *Front Microbiol* 3:110. <http://dx.doi.org/10.3389/fmicb.2012.00110>.
3. Naseer U, Sundsfjord A. 2011. The CTX-M conundrum: dissemination of plasmids and *Escherichia coli* clones. *Microb Drug Resist* 17:83–97. <http://dx.doi.org/10.1089/mdr.2010.0132>.
4. Cantón R, Coque TM. 2006. The CTX-M beta-lactamase pandemic. *Curr Opin Microbiol* 9:466–475. <http://dx.doi.org/10.1016/j.mib.2006.08.011>.

5. Luvsansharav UO, Hirai I, Niki M, Nakata A, Yoshinaga A, Moriyama T, Yamamoto Y. 2011. Prevalence of fecal carriage of extended-spectrum β -lactamase-producing *Enterobacteriaceae* among healthy adult people in Japan. *J Infect Chemother* 17:722–725. <http://dx.doi.org/10.1007/s10156-011-0225-2>.
6. Geser N, Stephan R, Korczak BM, Beutin L, Hächler H. 2012. Molecular identification of extended-spectrum- β -lactamase genes from *Enterobacteriaceae* isolated from healthy human carriers in Switzerland. *Antimicrob Agents Chemother* 56:1609–1612. <http://dx.doi.org/10.1128/AAC.05539-11>.
7. Zurfluh K, Hächler H, Stephan R, Nüesch-Inderbinen M. Long-term shedding of CTX-M-15 producing *E. coli* B2:ST127 by a healthy asymptomatic carrier. *Int J Antimicrob Agents*, in press.
8. Angiuoli SV, Gussman A, Klimke W, Cochrane G, Field D, Garrity G, Kodira CD, Kyrpides N, Madupu R, Markowitz V, Tatusova T, Thomson N, White O. 2008. Toward an online repository of Standard Operating Procedures (SOPs) for (meta)genomic annotation. *Omics* 12:137–141. <http://dx.doi.org/10.1089/omi.2008.0017>.
9. Alghoribi MF, Gibreel TM, Dodgson AR, Beatson SA, Upton M. 2014. *Galleria mellonella* infection model demonstrates high lethality of ST69 and ST127 uropathogenic *E. coli*. *PLoS One* 9:e101547. <http://dx.doi.org/10.1371/journal.pone.0101547>.
10. Gibreel TM, Dodgson AR, Cheesbrough J, Fox AJ, Bolton FJ, Upton M. 2012. Population structure, virulence potential and antibiotic susceptibility of uropathogenic *Escherichia coli* from Northwest England. *J Antimicrob Chemother* 67:346–356. <http://dx.doi.org/10.1093/jac/dkr451>.
11. Foxman B. 2010. The epidemiology of urinary tract infection. *Nat Rev Urol* 7:653–660. <http://dx.doi.org/10.1038/nrurol.2010.190>.